

Neutrino Mixing and Cosmological Constant above GUT Scale

Bipin Singh Koranga

November 3, 2009

Department of Physics, Kirori Mal college (University of Delhi,) Delhi-110007,
India

Abstract

Neutrino mixing lead to a non zero contribution to the cosmological constant. We consider non renormalization $1/M_x$ interaction term as a perturbation of the neutrino mass matrix. We find that for the degenerate neutrino mass spectrum. We assume that the neutrino masses and mixing arise through physics at a scale intermediate between Planck Scale and the electroweak scale. We also assume, above the electroweak breaking scale, neutrino masses are nearly degenerate and their mixing is bimaximal. Quantum gravitational (Planck scale)effects lead to an effective $SU(2)_L \times U(1)$ invariant dimension-5 Lagrangian involving neutrino and Higgs fields, which gives rise to additional terms in neutrino mass matrix. There additional term can be considered to be perturbation of the GUT scale bi-maximal neutrino mass matrix. We assume that the gravitational interaction is flavour blind and we study the neutrino mixing and cosmological constant due to physics above the GUT scale.

Keywords:Neutrino Mixing,Cosmological constant,GUT scale

1 Introduction

The problem of cosmological constant is currently one of the most challenging open issue in theortical physics and cosmology. The main difficulty comes from the mis match between theortical and accepted number. Cosmology constant may arise from neutrino mixing [1]. In this case of neutrinos, cosmological density reletated to the mixing and mass difference among the different generations. Phenomenological consequences of non-trivial condensate structure of the flavour vacuum have been studied for neutrino oscillations and Beta decay [2,3]. The nature of the cosmology constant Λ is one of the most intersting issues in modern theortical physics ans cosmology. Experimental data coming from observation indicates that not only Λ is different from zero, Λ also dominates

the universe dynamics driving an accelerated expansion [4,5]. In this paper, we study the neutrino mixing due to Planck scale and contribution to cosmological constant. In Section 2, we summarige the neutrino mixing due to Planck scale effects. In Section 3, we discuss the neutrino mixing and cosmological constant due to Planck scale effects. Section 4 is devoted to the conclusions.

2 Neutrino Oscillation Parameter due to Planck Scale Effects

The neutrino mass matrix is assumed to be generated by the see saw mechanism [6,7,8]. We assume that the dominant part of neutrino mass matrix arise due to GUT scale operators and the lead to bi-maximal mixing. The effective gravitational interaction of neutrino with Higgs field can be expressed as $SU(2)_L \times U(1)$ invarinat dimension-5 operator [8],

$$L_{grav} = \frac{\lambda_{\alpha\beta}}{M_{pl}} (\psi_{A\alpha} \epsilon \psi_C) C_{ab}^{-1} (\psi_{B\beta} \epsilon_{BD} \psi_D) + h.c. \quad (1)$$

Here and every where we use Greek indices α, β for the glavour states and Latin indices i, j, k for the mass states. In the above equation $\psi_\alpha = (\nu_\alpha, l_\alpha)$ is the lepton doublet, $\phi = (\phi^+, \phi^0)$ is the Higgs doublet and $M_{pl} = 1.2 \times 10^{19} GeV$ is the Planck mass λ is a 3×3 matrix in a flavour space with each elements $O(1)$. The Lorentz indices $a, b = 1, 2, 3, 4$ are contracted with the charge conjugation matrix C and the $SU(2)_L$ isospin indices $A, B, C, D = 1, 2$ are contracted with $\epsilon = i\sigma_2$, $\sigma_m (m = 1, 2, 3)$ are the pauli matrices. After spontaneous electroweak symmetry breaking the lagrangian in eq(1) generated additional term of neutrino mass matrix

$$L_{mass} = \frac{v^2}{M_{pl}} \lambda_{\alpha\beta} \nu_\alpha C^{-1} \nu_\beta, \quad (2)$$

where $v = 174 GeV$ is the VEV of electroweak symmetric breaking. We assume that the gravitational interaction is "flavour blind" that is $\lambda_{\alpha\beta}$ is independent of α, β indices. Thus the Planck scale contibution to the neutrino mass matrix is

$$\mu\lambda = \mu \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix}, \quad (3)$$

where the scale μ is

$$\mu = \frac{v^2}{M_{pl}} = 2.5 \times 10^{-6} eV. \quad (4)$$

We take eq(3) as perturbation to the main part of the neutrino mass matrix, that is generated by GUT dynamics. To calculate the effects of perturbation

on neutrino observables. The calculation developed in an earlier paper [8]. A natural assumption is that unperturbed (0^{th} order mass matrix) M is given by

$$\mathbf{M} = U^* \text{diag}(M_i) U^\dagger, \quad (5)$$

where, $U_{\alpha i}$ is the usual mixing matrix and M_i , the neutrino masses is generated by Grand unified theory. Most of the parameter related to neutrino oscillation are known, the major expectation is given by the mixing elements U_{e3} . We adopt the usual parametrization.

$$\frac{|U_{e2}|}{|U_{e1}|} = \tan\theta_{12}, \quad (6)$$

$$\frac{|U_{\mu 3}|}{|U_{\tau 3}|} = \tan\theta_{23}, \quad (7)$$

$$|U_{e3}| = \sin\theta_{13}. \quad (8)$$

In term of the above mixing angles, the mixing matrix is

$$U = \text{diag}(e^{if_1}, e^{if_2}, e^{if_3}) R(\theta_{23}) \Delta R(\theta_{13}) \Delta^* R(\theta_{12}) \text{diag}(e^{ia_1}, e^{ia_2}, 1). \quad (9)$$

The matrix $\Delta = \text{diag}(e^{\frac{i\delta}{2}}, 1, e^{-\frac{i\delta}{2}})$ contains the Dirac phase. This leads to CP violation in neutrino oscillation a_1 and a_2 are the so called Majoring phase, which effects the neutrino less double beta decay. f_1 , f_2 and f_3 are usually absorbed as a part of the definition of the charge lepton field. Planck scale effects will add other contribution to the mass matrix that gives the new mixing matrix can be written as [8]

$$U' = U(1 + i\delta\theta),$$

$$\begin{pmatrix} U_{e1} & U_{e2} & U_{e3} \\ U_{\mu 1} & U_{\mu 2} & U_{\mu 3} \\ U_{\tau 1} & U_{\tau 2} & U_{\tau 3} \end{pmatrix} + i \begin{pmatrix} U_{e2}\delta\theta_{12}^* + U_{e3}\delta\theta_{23}^*, & U_{e1}\delta\theta_{12} + U_{e3}\delta\theta_{23}^*, & U_{e1}\delta\theta_{13} + U_{e3}\delta\theta_{23}^* \\ U_{\mu 2}\delta\theta_{12}^* + U_{\mu 3}\delta\theta_{23}^*, & U_{\mu 1}\delta\theta_{12} + U_{\mu 3}\delta\theta_{23}^*, & U_{\mu 1}\delta\theta_{13} + U_{\mu 3}\delta\theta_{23}^* \\ U_{\tau 2}\delta\theta_{12}^* + U_{\tau 3}\delta\theta_{23}^*, & U_{\tau 1}\delta\theta_{12} + U_{\tau 3}\delta\theta_{23}^*, & U_{\tau 1}\delta\theta_{13} + U_{\tau 3}\delta\theta_{23}^* \end{pmatrix}. \quad (10)$$

Where $\delta\theta$ is a hermition matrix that is first order in μ [8,9]. The first order mass square difference $\Delta M_{ij}^2 = M_i^2 - M_j^2$, get modified [8,9] as

$$\Delta M_{ij}'^2 = \Delta M_{ij}^2 + 2(M_i \text{Re}(m_{ii}) - M_j \text{Re}(m_{jj})), \quad (11)$$

where

$$m = \mu U^t \lambda U,$$

$$\mu = \frac{v^2}{M_{pl}} = 2.5 \times 10^{-6} eV.$$

The change in the elements of the mixing matrix, which we parametrized by $\delta\theta$ [8,9], is given by

$$\delta\theta_{ij} = \frac{iRe(m_{jj})(M_i + M_j) - Im(m_{jj})(M_i - M_j)}{\Delta M_{ij}^2}. \quad (12)$$

The above equation determine only the off diagonal elements of matrix $\delta\theta_{ij}$. The diagonal element of $\delta\theta_{ij}$ can be set to zero by phase invariance. Using Eq(10), we can calculate neutrino mixing angle due to Planck scale effects,

$$\frac{|U'_{e2}|}{|U'_{e1}|} = \tan\theta'_{12}, \quad (13)$$

$$\frac{|U'_{\mu 3}|}{|U'_{\tau 3}|} = \tan\theta'_{23}, \quad (14)$$

$$|U'_{e3}| = \sin\theta'_{13} \quad (15)$$

For degenerate neutrinos, $M_3 - M_1 \cong M_3 - M_2 \gg M_2 - M_1$, because $\Delta_{31} \cong \Delta_{32} \gg \Delta_{21}$. Thus, from the above set of equations, we see that U'_{e1} and U'_{e2} are much larger than U'_{e3} , $U'_{\mu 3}$ and $U'_{\tau 3}$. Hence we can expect much larger change in θ_{12} compared to θ_{13} and θ_{23} [10]. As one can see from the above expression of mixing angle due to Planck scale effects, depends on new contribution of mixing $U' = U(1 + i\delta\theta)$.

3 Neutrino Mixing and Cosmological Constant Due to Planck Scale Effects

The connection between the vacuum energy density $\langle \rho_{vac} \rangle$ and the cosmology constant Λ is provided by the well known relation

$$\langle \rho_{vac} \rangle = \frac{\Lambda}{4\pi G}, \quad (16)$$

where G is the gravitational constant.

The expression of vacuum energy density $\langle \rho_{vac}^{mix} \rangle$ due to neutrino mixing is given by [11,12,13]

$$\langle \rho_{vac}^{mix} \rangle = 32\pi^2 \sin^2\theta_{12} \int dk K^2(\omega_{k,1} + \omega_{k,2}) |V_k|^2, \quad (17)$$

If we chose $K \gg \sqrt{m_1 m_2}$, we obtain

$$\langle \rho_{vac}^{mix} \rangle = \alpha \sin^2 \theta_{12} (m_2 - m_1)^2 K^2 \int_0^k dk k^2 (\omega_{k,1} + \omega_{k,2}) |V_K|^2, \quad (18)$$

For hierarchical neutrino model, for which $m_2 > m_1$, we have in this case $K \gg \sqrt{m_1 m_2}$ and take into account the asymptotic properties of V_k

$$|V_k|^2 \simeq \frac{(m_2 - m_1)^2}{4K^2}, \quad K \gg \sqrt{m_1 m_2}$$

We get

$$\langle \rho_{vac}^{mix} \rangle = \alpha \sin^2 \theta_{12} (m_2^2 - m_1^2) \propto \frac{\Lambda}{4\pi G}, \quad (19)$$

The new cosmological constant Λ due to Planck scale effects is given by

$$\Lambda' = \alpha \sin^2 \theta'_{12} (m_2^2 - m_1^2), \quad (20)$$

where θ'_{12} is given by eq(13)

We consider the Planck scale effects on neutrino mixing and we get the given range of mixing parameter of MNS matrix

$$U' = R(\theta_{23} + \epsilon_3) U_{phase}(\delta) R(\theta_{13} + \epsilon_2) R(\theta_{12} + \epsilon_1). \quad (21)$$

In Planck scale, only $\theta_{12}(\epsilon_1 = \pm 3^\circ)$ have reasonable deviation and θ_{23}, θ_{13} deviation is very small less than 0.3° [10]. In the new mixing at Planck scale we get the cosmological density

$$\Lambda' = \alpha \sin^2 (\theta_{12} \pm \epsilon_1) (m_2'^2 - m_1^2), \quad (22)$$

The presence of a cosmological constant fluid has to be compatible with the structure formation, allow to set the upper bound $\Lambda < 10^{-56} \text{cm}^{-2}$ [14]. Due to Planck scale effects mixing angle θ_{12} deviated the cosmological constant Λ .

4 Conclusions

We assume that the main part of neutrino masses and mixing from GUT scale operator. We considered these to be 0^{th} order quantities. We further assume that GUT scale symmetry constrain the neutrino mixing angles to be bimaximal. The gravitational interaction of lepton field with S.M Higgs field give rise to a $SU(2)_L \times U(1)$ invariant dimension-5 effective lagrangian give originally by Weinberg [15]. On electroweak symmetry breaking this operators leads to additional mass terms. We considered these to be perturbation of GUT scale mass terms. We compute the first order correction to neutrino mass eigen value

and mixing angles. In [10], it was shown that the change in θ_{13}, θ_{23} is very small (less than 0.3°) but the change in θ_{12} can be substantial about $\pm 3^\circ$. The change in all the mixing angles are proportional to the neutrino mass eigenvalues. To maximize the change, we assumed degenerate neutrino mass 2.0eV . For degenerate neutrino masses, the change in θ_{13}, θ_{23} are inversely proportional to Δ_{21} . Since $\Delta_{31} \cong \Delta_{32} \gg \Delta_{21}$, the change in θ_{12} is much larger than the change in other mixing angles. In this paper, we write the cosmological constant above GUT scale in terms of mixing angles for Majorana neutrinos, these expressions are in eq.(x) for vacuum mixing. For Majorana neutrinos, the expression is $\Lambda \propto \sin^2 \theta_{12} (m_2^2 - m_1^2)$. In this paper, finally we wish to make an important comment. Due to Planck scale effects, the mixing angle θ_{12} deviates from the cosmological constant Λ .

References

- [1] M. Blasone, *et al.*, Phys. Lett. **A323**, 182 (2004).
- [2] M. Blasone, *et al.*, Phys. Rev. **D67**, 073011 (2003).
- [3] M. Blasone, *et al.*, hep-ph/0307205.
- [4] S. Perlmutter *et al.*, Astrophys. J. **517**, 565 (1999).
- [5] V. Sahni and A. Starobinsky, Int. J. Mod. Phys. **D9**, 373 (2000).
- [6] R.N. Mohapatra *et al.*, Phys. Rev. Lett. **44**, 912 (1980).
- [7] S. Coleman and S. L. Glashow, Phys. Rev. **D59**, 116008 (1999).
- [8] F. Vissani *et al.*, Phys. Lett. **B571**, 209, (2003).
- [9] Bipin Singh Koranga, Mohan Narayan and S. Uma Sankar, arXiv:hep-ph/0611186.
- [10] Bipin Singh Koranga, Mohan Narayan and S. Uma Sankar, Phys. Lett. **B665**, 63 (2008).
- [11] M. Blasone, *et al.*, Phys. Lett. **A323**, 182 (2004).
- [12] M. Blasone, *et al.*, Braz. J. Phys. **35**:447-454 (2005).
- [13] A. Capolupo, S. Capozziello and G. Vitiello, Phys. Lett. **A363**, 53 (2007).
- [14] Ya.B. Zeldovich, I.D. Novikov, Structure and evolution of the universe Moscow, Izdatelstvo Nauka (1975).
- [15] S. Weinberg, Phys. Rev. Lett. **43**, 1566 (1979).